

Baldwin Hills Air Quality Study

Michael McCarthy, Steven Brown, Song Bai, Jennifer DeWinter,
Theresa O'Brien, David Vaughn, Paul Roberts
Sonoma Technology, Inc., Petaluma, CA

Richard Peltier
University of Massachusetts at Amherst, Environmental Health Sciences

Shane Murphy
University of Wyoming, Atmospheric Science

for

National Ambient Air Monitoring Conference
Atlanta, GA

August 13, 2014




Sonoma Technology, Inc.

Outline

- Background and objectives
- Methods
- Results
- Summary

Motivation

- Multiple community groups were concerned with potential pollutant impacts from oil field activities.
- Conducted a 1-year study as part of a settlement about an Environmental Impact Report (EIR) on oil field development.



Aerial view of the Inglewood Oil Field, showing the locations of the four monitoring sites: north (N), east (E), south (S), and west (W).

Objectives ⁽¹⁾

Primary project objectives


- Quantify air toxics emissions from Inglewood Oil Field operations.
- Assess the health risk of acute and chronic exposure to air toxics emitted from oil field operations.



Objectives ⁽²⁾

Secondary project objectives (to the extent feasible)

- Determine and distinguish major sources of toxic air emissions within the areas surrounding Inglewood Oil Field.
- Assess the oil field's contribution to the overall acute and chronic health risk in the surrounding areas.



Challenges

- The oil field is in an urban area of 15 million people with other major sources (refineries, LAX, etc.) in the area.
- Houses have been built right up to the edge of the oil field.
- A regular diurnal wind pattern puts different parts of the nearby community downwind of the oil field during different parts of the day.
- It would be very expensive to monitor all air toxics at all impacted locations.

Prioritized Toxics of Concern

Key pollutants identified for characterization

- Diesel particulate matter (DPM)
- Cadmium, nickel, mercury, manganese, arsenic, lead
- Benzene, formaldehyde, acrolein


Addressing Challenges

- Collect high time-resolution data; daily or greater average data will not be as useful because wind patterns shift multiple times a day, making apportionment difficult.
- Select monitoring sites to
 - Represent highest and longest potential impact from the oil field
 - Determine the incremental change in ambient concentrations across the oil field
- All within a limited budget.

Monitoring Methods

- **Black carbon** (BC; as a proxy for DPM) by Teledyne-API Model 633 dual-wavelength Aethalometers.
- **24 metals** by XACT 625 semi-continuous X-ray fluorescence (XRF) spectrometer (e.g., S, K, Fe, Cu, Pb, Ni, Cd, As, Hg).
- **7 VOCs** by Proton-Transfer-Reaction Time-of-Flight Mass Spectrometry (PTR-TOFMS) (e.g., acrolein, benzene, butadiene, naphthalene, acetaldehyde).
- **Meteorology** for wind patterns, temperature, and humidity.

- Hourly BC at all sites 11/15/12-11/15/13
- Hourly metals 11/15/12-2/1/13
- 1-min VOCs 7/3/13-7/17/13
- Meteorology 11/15/12-11/15/13




Results

- Monitoring was completed in November 2013 and data analyzed to characterize
 - Temporal patterns of pollutants
 - Relative contribution of sources to pollutants
 - Hazard associated with oil field operations
- Currently working with Los Angeles County on follow-up analyses linking ambient concentrations with nearby vehicle activity and facility operations.


BC Concentrations

- BC was typically below $1 \mu\text{g}/\text{m}^3$ at all sites.
- Morning peak in BC.
- Operations on oil field were 24/7/365, but drilling was limited to daytime only.





BC Differential

- Predominant west-southwest wind direction (53% of hours).
- We can use the difference between the S and E sites to quantify the contribution from the oil field area.




BC Differential

- Daytime concentrations of BC were consistently higher at the downwind site during daytime working hours.
- Overnight, when there were no operations, differential was typically zero.
- The differential indicates
 - Statistically significant BC from oil field operations
 - But with a contribution of $\sim 0.1 \mu\text{g}/\text{m}^3$





XRF Metals

- Temporal & wind direction patterns were examined
- EPA PMF was used to evaluate potential sources




Metals

- Cu, Mn, and Pb show daytime spikes, similar to BC differential; other species showed mixed patterns.
- No species showed significantly higher concentrations when site was downwind of the oil field.




Metals Apportionment

- Identified multiple sources, including a factor with high Mn and Ni, likely associated with oil field operations.
- Spikes in Mn/Ni factor occurred coincident with operations activity and higher BC differential.



VOCs

- VOCs generally showed a similar diurnal pattern as BC.
- No consistent signal from oil operations was found.
- Some transient high concentration events occurred coincident with operations, but had low influence on average.



Summary

- Focus resources on the most important issues—in this case, DPM drives the risk estimates.
- Balance scope of work with available budget; year-round monitoring of metals, BC, and VOCs would be expensive, and may not be of sufficient temporal resolution.
- While transient high concentration events occurred coincident with oil field operations, there was no clear influence on ambient metals and VOCs.
- Only a small fraction of BC was attributed to oil field operations.

Reference & Acknowledgments

- California Environmental Protection Agency
<http://www.oehha.ca.gov/air/allrels.html>.
- Funding provided by the Los Angeles County Department of Regional Planning.

Contacts



Steve Brown

Sr. Atmospheric Scientist

steveb@sonomatech.com



Hilary Hafner

Sr. Vice President

hilary@sonomatech.com



Sonoma Technology, Inc.

Air Quality Research and Innovative Solutions

sonomatech.com

[@sonoma_tech](https://twitter.com/sonoma_tech)